

Tilburg University

Uniform approximations of the stationary and palm distributions of marked point processes

Nieuwenhuis, G.

Publication date:
1992

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Nieuwenhuis, G. (1992). *Uniform approximations of the stationary and palm distributions of marked point processes*. (Research Memorandum FEW). Faculteit der Economische Wetenschappen.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ECBM

R

7626

1992

543

UNIVERSITY
LEIJE
UNIVERSITEIT
BRABANT

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS



DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM

UNIFORM APPROXIMATIONS OF THE
STATIONARY AND PALM DISTRIBUTIONS OF
MARKED POINT PROCESSES


Gert Nieuwenhuis

R 43

FEW 543

*Stationary Point
Approximation Theory*

Refereed by Prof.dr. B.B. van der Genugten



K.U.B.
BIBLIOTHEEK
TILBURG

Uniform Approximations of the Stationary and Palm Distributions of Marked Point Processes

Gert Nieuwenhuis
Tilburg University
Department of Econometrics
P.O. Box 90153
NL-5000 LE Tilburg
The Netherlands

Summary

Let P be the distribution of a stationary marked point process on \mathbf{R} and let P_L^0 be its Palm distribution with respect to a set L of marks. The probability measures $P_{i,L}$, $i \in \mathbf{Z}$, arise from P by shifting the origin to the i th occurrence with mark in L . In Nieuwenhuis (1991) the well-known approximation of P_L^0 by the mean of $P_{1,L}, \dots, P_{n,L}$ under an ergodicity condition was proved to be uniform. If this condition is not satisfied, then the (uniform) limit of this mean can still be characterized.

In this paper it is proved that in the results just mentioned P may be replaced by $P_{L'}^0$, where L' is another set of marks with $L \cap L' = \emptyset$. In a 'dual' theorem the roles of P and P_L^0 are interchanged. Starting from P_L^0 the uniform convergence of a Césaro mean of shifted probabilities is considered. Under a weak ergodicity condition the limit is equal to P .

AMS 1980 subject classifications. Primary 60G55; secondary 60G10, 60F15.

Key words and phrases. Palm distribution, marked point process, cross-convergence results.

1 Introduction

Let K be a complete and separable metric space. A *marked point process on \mathbf{R} with mark space K* is a random element Φ in the class of all integer-valued measures φ on the σ -field $\text{Bor } \mathbf{R} \times \text{Bor } K$ such that:

$$\varphi(A \times K) < \infty \text{ for all bounded } A \in \text{Bor } \mathbf{R}.$$

Let M_K be this class and endow it with the σ -field \mathcal{M}_K generated by the sets $[\varphi(A \times L) = k] := \{\varphi \in M_K : \varphi(A \times L) = k\}$, $k \in \mathbf{N}_0$, $L \in \text{Bor } K$ and $A \in \text{Bor } \mathbf{R}$. The distribution of Φ will be denoted by P , a probability measure on (M_K, \mathcal{M}_K) .

For $\varphi \in M_K$ and $L \in \text{Bor } K$ we define $\tilde{\varphi}_L \in M_K$ and the counting measure φ_L on $\text{Bor } \mathbf{R}$ by $\tilde{\varphi}_L(B) := \varphi(B \cap (\mathbf{R} \times L))$ and $\varphi_L(A) := \varphi(A \times L)$, $B \in \text{Bor } \mathbf{R} \times \text{Bor } K$ and $A \in \text{Bor } \mathbf{R}$. Set

$$\begin{aligned} M_L^\infty &:= \{\varphi \in M_K : \varphi_L(-\infty, 0) = \varphi_L(0, \infty); \varphi_K(\{s\}) \leq 1 \text{ for all } s \in \mathbf{R}\}, \\ M_L^0 &:= \{\varphi \in M_L^\infty : \varphi_L(\{0\}) = 1\}, \\ \mathcal{M}_L^\infty &:= M_L^\infty \cap \mathcal{M}_K \text{ and } \mathcal{M}_L^0 := M_L^0 \cap \mathcal{M}_K, \end{aligned}$$

$L \in \text{Bor } K$. Define $\lambda(L) := \mathbf{E}\Phi_L(0, 1]$. It will always be assumed that Φ (or rather P) is *stationary* (i.e., $\Phi(t + \cdot) =_d \Phi$ for all $t \in \mathbf{R}$), that $P(M_K^\infty) = 1$, and that the *intensity* $\lambda(K)$ is finite. We will only consider $L \in \text{Bor } K$ with $P(M_L^\infty) = 1$. The atoms of $\varphi \in M_K^\infty$ are denoted by $(X_i(\varphi), k_i(\varphi)) \in \mathbf{R} \times K$, $i \in \mathbf{Z}$, with the convention

$$\dots < X_{-1}(\varphi) < X_0(\varphi) \leq 0 < X_1(\varphi) < \dots$$

$X_i(\varphi)$ is interpreted as the i th *occurrence* (or *point*) of φ , $k_i(\varphi)$ as the accessory *mark*. For $\varphi \in M_L^\infty$ we write $X_i^L(\varphi) := X_i(\tilde{\varphi}_L)$, the ' i th L -point of φ ', and $\alpha_i^L(\varphi) := X_{i+1}^L(\varphi) - X_i^L(\varphi)$.

Two types of shifts will be considered. The *time shifts* $T_t : M_K^\infty \rightarrow M_K^\infty$, $t \in \mathbf{R}$, are determined by $T_t\varphi(A \times L) := \varphi((t + A) \times L)$, $\varphi \in M_K^\infty$, $A \in \text{Bor } \mathbf{R}$ and $L \in \text{Bor } K$. Note that $\varphi(t + \cdot) = T_t\varphi$ can be represented by $\{(X_i(\varphi) - t, k_i(\varphi)) : i \in \mathbf{Z}\}$. For fixed $L \in \text{Bor } K$ with $P(M_L^\infty) = 1$ the *point shifts* $\vartheta_{n,L} : M_L^\infty \rightarrow M_L^\infty$, $n \in \mathbf{Z}$, are defined by

$\vartheta_{n,L}\varphi := \varphi(X_n^L(\varphi) + \cdot)$. The probability measures $P_{n,L} := P\vartheta_{n,L}^{-1}$, $n \in \mathbb{Z}$, on $(M_L^\infty, \mathcal{M}_L^\infty)$ arise from P by shifting the origin to the n th occurrence.

Let $L \in \text{Bor } K$ be such that $P(M_L^\infty) = 1$. We now consider the Palm distribution P_L^0 of Φ with respect to L . The formal definition of P_L^0 follows below, but intuitively P_L^0 is the conditional distribution of Φ given the occurrence of an arbitrary L -point in the origin. This intuitive meaning of P_L^0 can (at least in the ergodic case) indeed be confirmed, since $P_L^0(B)$, $B \in \mathcal{M}_L^\infty$, can be approximated by

$$\frac{1}{n} \sum_{i=1}^n P[\vartheta_{n,L}\varphi \in B], \quad n \in \mathbb{N}, \quad (1.1)$$

if Φ satisfies some weak ergodicity condition. Here $[\vartheta_{n,L}\varphi \in B] := \{\varphi \in M_L^\infty : \vartheta_{n,L}\varphi \in B\}$. In Nieuwenhuis (1991) it has been proved that the convergence in question holds uniformly for $B \in \mathcal{M}_L^\infty$. If, however, this ergodicity condition is not satisfied, then the (uniform) limit $Q_L^0(B)$ of the sequence in (1.1) is not necessarily equal to $P_L^0(B)$, but can still be characterized. See Theorem 1.2 below.

In Section 4 it is proved that in the above results (as stated in Theorem 1.2) P may be replaced by $P_{L'}^0$, where $L' \in \text{Bor } K$ is another set of marks, $L \cap L' = \emptyset$. The resulting theorem concerns the limit behavior of $(n^{-1} \sum_{i=1}^n P_{L'}^0[\vartheta_{n,L}\varphi \in B])_{n \in \mathbb{N}}$. It can be compared with Theorem 3 in Konstantopoulos and Walrand (1988), which in essence concerns weak convergence of $(P_{L'}^0[\vartheta_{n,L}\varphi \in \cdot])_{n \in \mathbb{N}}$ under some mixing condition. See also König and Schmidt (1986).

In Section 3 the roles of P and P_L^0 in Theorem 1.2 are interchanged. It is proved that the Césaro mean

$$\frac{1}{t} \int_0^t P_L^0[T_x\varphi \in B] dx, \quad t \in (0, \infty), \quad (1.2)$$

tends to some limit $Q_L(B)$ as $t \rightarrow \infty$, uniformly for $B \in \mathcal{M}_L^\infty$. The probability measure Q_L turns out to be equal to P if some weak ergodicity condition holds. The probability measures Q_L and Q_L^0 are connected.

Our treatment involves conditioning on invariant σ -fields. Some preliminary lemmas are proved in Section 2.

In our proves we have to go from P_L^0 to P or from P to P_L^0 , several times. The method used to bridge these gaps (the ‘Radon-Nikodym approach’, see Nieuwenhuis (1991; Section 1)), is a consequence of Theorem 1.1.

We next formalize some of the notions mentioned above and give some other definitions and notations.

For $L \in \text{Bor } K$ with $P(M_L^\infty) = 1$ the *Palm distribution* P_L^0 of Φ (or rather P) with respect to L is defined by

$$P_L^0(A) := \frac{1}{\lambda(L)} \mathbf{E} \left[\sum_{i=1}^{\Phi((0,1] \times L)} 1_A(\vartheta_{i,L} \Phi) \right], \quad A \in \mathcal{M}_L^\infty. \quad (1.3)$$

Intuitively P_L^0 arises from P by shifting the origin to an arbitrary L -point. Note the difference between P_L^0 and $P_{0,L}$, in notation as well as in interpretation. Several probability measures on $(M_L^\infty, \mathcal{M}_L^\infty)$ have been defined now: P , P_L^0 , $P_{n,L}$. In this research expectations with respect to these measures are denoted by E , E_L^0 , $E_{n,L}$, respectively. When another probability measure Q on $(M_L^\infty, \mathcal{M}_L^\infty)$ is considered, we will write E_Q . Expectation with respect to an universal probability space (Ω, \mathcal{F}, P) is (as in (1.3)) denoted by \mathbf{E} .

Note that $P_L^0(M_L^0) = 1$. The probability measure P_L^0 has the following properties:

$$P_L^0 \vartheta_{n,L}^{-1} = P_L^0 \quad \text{for all } n \in \mathbb{Z}, \quad (1.4)$$

$$P(A) = \lambda(L) \int_0^\infty P_L^0[X_1^L(\varphi) > u; \varphi(u + \cdot) \in A] du, \quad A \in \mathcal{M}_L^\infty. \quad (1.5)$$

See Franken et al. (1982), Matthes, Kerstan and Mecke (1978), Kallenberg (1983/86), and Brandt, Franken and Lisek (1990) for more information.

The *inversion formula* (1.5) expresses P in terms of P_L^0 ; the definition in (1.3) expresses P_L^0 in terms of P . There is another way of going from P_L^0 to P (and vice versa). The essence of the approach in question is contained in the next theorem. It is proved in Nieuwenhuis (1989), with an extension to **marked** point processes in Nieuwenhuis (1991; Section 5).

(Two probability distributions Q_1 and Q_2 on a common measurable space are said to be equivalent (notation $Q_1 \sim Q_2$) if they have the same null-sets. A Radon-Nikodym derivative of Q_1 with respect to Q_2 is denoted by $\frac{dQ_1}{dQ_2}$.)

Theorem 1.1 *Let $n \in \mathbb{Z}$ and let $L \in \text{Bor } K$ be such that $P(M_L^\infty) = 1$. Then*

- (i) $P_{n,L} \sim P_L^0$,
- (ii) $\frac{dP_{n,L}}{dP_L^0} = \lambda(L)\alpha_{-n}^L \quad P_L^0\text{-a.s.}$

Suppose that $f : M_L^0 \rightarrow \mathbb{R}$ is P_L^0 -integrable. We are now able to express the P_L^0 -expectation of f in terms of a P -expectation:

$$E_L^0 f = \frac{1}{\lambda(L)} E_{0,L} \left(\frac{1}{\alpha_0^L} f \right) = \frac{1}{\lambda(L)} E \left(\frac{1}{\alpha_0^L} f \circ \vartheta_{0,L} \right). \quad (1.6)$$

Reversely, if $g : M_L^\infty \rightarrow \mathbb{R}$ is P -integrable with $Eg = Eg \circ \vartheta_{0,L}$, then P -expectation of g can be transformed to a P_L^0 -expectation:

$$Eg = E_{0,L}g = \lambda(L)E_L^0(\alpha_0^L g). \quad (1.7)$$

For more information we refer to Nieuwenhuis (1991). The approach in (1.6) and (1.7), where $P_{0,L}$ is used as a bridge between P_L^0 and P , is very common in this research.

Consider the following invariant σ -fields:

$$\begin{aligned} \mathcal{I}'_L &:= \{A \in \mathcal{M}_L^\infty : T_t^{-1}A = A \text{ for all } t \in \mathbb{R}\} \text{ and} \\ \mathcal{I}_L &:= \{A \in \mathcal{M}_L^\infty : \vartheta_{1,L}^{-1}A = A\}. \end{aligned} \quad (1.8)$$

Φ (or rather P) is called *ergodic* if $P(A) \in \{0, 1\}$ for all $A \in \mathcal{I}'_K$; it is called *pseudo- L -ergodic* if

$$E_L^0(\alpha_0^L | \mathcal{I}_L) = \frac{1}{\lambda(L)} \quad P_L^0\text{-a.s.} \quad (1.9)$$

P_L^0 is called *ergodic* if $P_L^0(A) \in \{0, 1\}$ for all $A \in \mathcal{I}_L$.

The following theorem has been the inspiration and motivation for this research. In this *cross-convergence* result P_L^0 is approximated when starting from P . It is proved in Nieuwenhuis (1991; Sections 4 and 5).

Theorem 1.2 *Let $L \in \text{Bor } K$ be such that $P(M_L^\infty) = 1$. Then*

$$\frac{1}{n} \sum_{i=1}^n P[\vartheta_{i,L} \varphi \in B] \rightarrow E(E_L^0(1_B | \mathcal{I}_L)) =: Q_L^0(B) \quad (1.10)$$

uniformly for $B \in \mathcal{M}_L^\infty$. The probability measure Q_L^0 on $(M_L^\infty, \mathcal{M}_L^\infty)$ is equivalent to P_L^0 and

$$\frac{dQ_L^0}{dP_L^0} = \lambda(L) E_L^0(\alpha_0^L | \mathcal{I}_L) \quad P_L^0 - a.s.$$

Q_L^0 and P_L^0 are equal iff Φ is pseudo- L -ergodic.

Let Q_1 and Q_2 be probability measures on a common measurable space, both dominated by a σ -finite measure μ and having densities h_1 and h_2 respectively. The *total variation distance* between Q_1 and Q_2 is defined by

$$d(Q_1, Q_2) := \int |h_1 - h_2| d\mu.$$

It is well-known that

$$d(Q_1, Q_2) = 2 \sup_A |Q_1(A) - Q_2(A)| = 2(Q_1[h_1 \geq h_2] - Q_2[h_1 \geq h_2]). \quad (1.11)$$

As a final remark we note that, when talking about Radon-Nikodyn derivatives, the supplement a.s. (almost surely) is often suppressed.

2 Conditioning on invariant σ -fields

One of the objectives of this research is to obtain some uniform cross- convergence results without assuming ergodicity. To realize this in this general setting we will condition on invariant σ -fields.

Recall the definitions of \mathcal{I}_L and \mathcal{I}_L' in (1.8).

Lemma 2.1 *Let $L \in \text{Bor } K$. Then:*

- (a) If $A \in \mathcal{I}_L$, then $\vartheta_{i,L}^{-1}A = A$ for all $i \in \mathbb{Z}$.
- (b) $\mathcal{I}_L = \mathcal{I}'_L$.

Proof. Let $A \in \mathcal{I}_L$. For $\psi \in M_L^\infty$ we have

$$\psi \in A \quad \text{iff} \quad \vartheta_{1,L}\psi \in A. \quad (2.1)$$

Let $\varphi \in M_L^\infty$. Since $\vartheta_{1,L} = \vartheta_{1,L} \circ \vartheta_{0,L}$, we obtain (by applying (2.1) twice) that

$$\varphi \in A \quad \text{iff} \quad \vartheta_{1,L}(\vartheta_{0,L}\varphi) \in A \quad \text{iff} \quad \vartheta_{0,L}\varphi \in A.$$

Consequently, $\vartheta_{0,L}^{-1}A = A$.

Since $\vartheta_{0,L} = \vartheta_{1,L} \circ \vartheta_{-1,L}$, we have for $\varphi \in M_L^\infty$, as another consequence of (2.1),

$$\varphi \in A \quad \text{iff} \quad \vartheta_{1,L}(\vartheta_{-1,L}\varphi) \in A \quad \text{iff} \quad \vartheta_{-1,L}\varphi \in A.$$

Hence, $\vartheta_{-1,L}^{-1}A = A$. Since $\vartheta_{i,L} = \vartheta_{1,L}^i$ and $\vartheta_{-i,L} = \vartheta_{-1,L}^i$ for all $i \geq 1$, part (a) follows immediately.

For $A \in \mathcal{I}'_L$ we have: $\varphi \in A$ iff $T_t\varphi \in A$, for all $t \in \mathbb{R}$ and $\varphi \in M_L^\infty$. Consequently, $\varphi \in A$ iff $\vartheta_{1,L}\varphi \in A$, for all $\varphi \in M_L^\infty$. So, $\vartheta_{1,L}^{-1}A = A$ and $\mathcal{I}'_L \subset \mathcal{I}_L$.

Let $A \in \mathcal{I}_L$. By (a) we have

$$\varphi \in A \quad \text{iff} \quad \vartheta_{i,L}\varphi \in A, \quad \text{for all } \varphi \in M_L^\infty \text{ and } i \in \mathbb{Z}. \quad (2.2)$$

Let $\varphi \in M_L^\infty$ and $t \in \mathbb{R}$. Take $i \in \mathbb{Z}$ such that $X_{i-1}^L(\varphi) \leq t < X_i^L(\varphi)$. Then

$$T_t\varphi \in A \quad \text{iff} \quad \vartheta_{1,L}(T_t\varphi) \in A \quad \text{iff} \quad \vartheta_{i,L}\varphi \in A \quad \text{iff} \quad \varphi \in A,$$

cf. (2.2). So, $T_t^{-1}A = A$ and $\mathcal{I}_L \subset \mathcal{I}'_L$. Part (b) follows immediately. \square

Note that as a consequence of Lemma 2.1 every \mathcal{I}_L -measurable function $f : M_L^\infty \rightarrow [0, \infty)$ satisfies

$$f \circ \vartheta_{i,L}(\varphi) = f(\varphi) \quad \text{and} \quad f \circ T_t(\varphi) = f(\varphi) \quad (2.3)$$

for all $\varphi \in M_L^\infty$, $i \in \mathbb{Z}$, and $t \in \mathbb{R}$.

In view of Section 4 we next consider two disjoint sets of marks. So, let $L, L' \in \text{Bor } K$ and $L \cap L' = \emptyset$. Furthermore, set

$$\begin{aligned} M_{L,L'}^\infty &:= M_L^\infty \cap M_{L'}^\infty \text{ and } \mathcal{M}_{L,L'}^\infty := M_{L,L'}^\infty \cap \mathcal{M}_K, \\ \mathcal{I}_{L,L'} &:= \{A \in \mathcal{M}_{L,L'}^\infty : \vartheta_{i,L}^{-1} A = A\}, \\ \mathcal{I}'_{L,L'} &:= \{A \in \mathcal{M}_{L,L'}^\infty : T_t^{-1} A = A \text{ for all } t \in \mathbb{R}\}. \end{aligned}$$

In this context the maps $\vartheta_{i,L}$ and T_t , $t \in \mathbb{R}$, will always be restricted to $M_{L,L'}^\infty$. The following relations can easily be proved:

$$\begin{aligned} \mathcal{I}'_L \cap M_{L'}^\infty &= \mathcal{I}'_{L,L'} \quad \text{and} \quad \mathcal{I}_L \cap M_{L'}^\infty = \mathcal{I}_{L,L'}; \\ \mathcal{I}'_{L,L'} &\subset \mathcal{I}'_L \quad \text{and} \quad \mathcal{I}_{L,L'} \subset \mathcal{I}_L \end{aligned} \quad (2.4)$$

At first sight part (b) of the next lemma seems rather surprising.

Lemma 2.2 *Let $L, L' \in \text{Bor } K$ with $L \cap L' = \emptyset$. Then:*

- (a) If $A \in \mathcal{I}_{L,L'}$, then $\vartheta_{i,L}^{-1} A = A$ for all $i \in \mathbb{Z}$;
- (b) $\mathcal{I}'_{L,L'} = \mathcal{I}_{L,L'} = \mathcal{I}'_{L',L}$.

Proof. The proof of (a) is similar to the proof of Lemma 2.1(a). Part (b) is an immediate consequence of Lemma 2.1(b) and (2.4) since

$$\begin{aligned} \mathcal{I}_{L,L'} &= \mathcal{I}_L \cap M_{L'}^\infty = \mathcal{I}'_L \cap M_{L'}^\infty = \mathcal{I}'_{L,L'} = \mathcal{I}'_{L',L} \\ &= \mathcal{I}'_{L'} \cap M_L^\infty = \mathcal{I}_{L',L} \cap M_L^\infty = \mathcal{I}_{L',L}. \end{aligned}$$

□

Next a stationary point process Φ with distribution P is put upon the stage. Suppose that $P(M_L^\infty) = 1$. Since $\mathcal{I}'_L \subset \mathcal{I}'_K$ and $\mathcal{I}'_L = \mathcal{I}'_K \cap \mathcal{M}_L^\infty$, the σ -field \mathcal{I}'_K in the definition

of ergodicity of P in Section 1 may equivalently be replaced by \mathcal{I}'_L . As a consequence of Lemma 2.1(b) we obtain:

$$\begin{aligned} P \text{ is ergodic} &\iff P_L^0 \text{ is ergodic,} \\ P \text{ is ergodic} &\implies P \text{ is pseudo-}L\text{-ergodic.} \end{aligned} \tag{2.5}$$

See also Nieuwenhuis (1991; Section 4).

In the following lemma some special conditional expectations are compared. The random variable $N_1^L : M_L^\infty \rightarrow \mathbf{N}_0$ is defined by $N_1^L(\varphi) := \varphi_L(0, 1]$.

Lemma 2.3 *Let $L, L' \in \text{Bor } K$ with $L \cap L' = \emptyset$ and $P(M_{L,L'}^\infty) = 1$. Then:*

- (a) $E_L^0(\alpha_0^L | \mathcal{I}_L) > 0$ and $E(\frac{1}{\alpha_0^L} | \mathcal{I}_L) = E(N_1^L | \mathcal{I}_L) = \frac{1}{E_L^0(\alpha_0^L | \mathcal{I}_L)} \quad P\text{- and } P_L^0\text{-a.s.}$
- (b) $E(\frac{1}{\alpha_0^L} | \mathcal{I}_{L,L'}) = \frac{1}{E_L^0(\alpha_0^L | \mathcal{I}_{L,L'})} \quad P_{L'}^0\text{-a.s.}$

Proof. Let $A \in \mathcal{I}_L$. Note that $\alpha_0^L = \alpha_0^L \circ \vartheta_{0,L}$. By (2.3), Theorem 1.1, and (1.3) we have

$$E(1_A E(\frac{1}{\alpha_0^L} | \mathcal{I}_L)) = E(1_A \frac{1}{\alpha_0^L}) = E_{0,L}(1_A \frac{1}{\alpha_0^L}) = \lambda(L) P_L^0(A) = E(1_A N_1^L).$$

Under P the first equality in the right-hand part of (a) is a consequence of this observation. Since P_L^0 and $P_{0,L}$ have the same null-sets, this equality is also valid under P_L^0 . (The left-hand part of (2.3) is used here.) Set $B := [E_L^0(\alpha_0^L | \mathcal{I}_L) \leq 0]$. Then

$$0 \geq E_L^0(1_B E_L^0(\alpha_0^L | \mathcal{I}_L)) = E_L^0(1_B \alpha_0^L).$$

Since $P_L^0[\alpha_0^L > 0] = 1$, we obtain

$$P_L^0(B^c) = 1 \quad \text{and} \quad P(B^c) = E(1_{B^c} \circ \vartheta_{0,L}) = P_{0,L}(B^c) = 1.$$

The left-hand part of (a) follows.

Let again $A \in \mathcal{I}_L$. By (2.3) and Theorem 1.1 we have

$$\begin{aligned}
E\left(1_A \frac{1}{E_L^0(\alpha_0^L|\mathcal{I}_L)}\right) &= E\left(1_A \circ \vartheta_{0,L} \frac{1}{E_L^0(\alpha_0^L|\mathcal{I}_L) \circ \vartheta_{0,L}}\right) \\
&= \lambda(L) E_L^0\left(\alpha_0^L 1_A \frac{1}{E_L^0(\alpha_0^L|\mathcal{I}_L)}\right) \\
&= \lambda(L) P_L^0(A) = E\left(1_A \frac{1}{\alpha_0^L}\right).
\end{aligned}$$

In the third equality we conditioned on \mathcal{I}_L . The right-hand part of (a) follows immediately.

Next (b). Since $\mathcal{I}_{L,L'} = \mathcal{I}_L \cap M_{L'}^\infty$ and $P(M_{L,L'}^\infty) = 1$, it is obvious that (a) remains valid if \mathcal{I}_L is replaced by $\mathcal{I}_{L,L'}$. So, part (b) holds under P . Since $\mathcal{I}_{L,L'} = \mathcal{I}_{L',L} \subset \mathcal{I}_{L'}$, both conditional expectations in (b) are $\mathcal{I}_{L'}$ -measurable. Hence, equality holds $P_{0,L'}$ -a.s. as well. By Theorem 1.1(ii) part (b) follows. \square

The following equality is proved in Nieuwenhuis (1991).

$$E_L^0\left(\int_0^{\alpha_0^L} g \circ T_s ds | \mathcal{I}_L\right) = E(g|\mathcal{I}_L) E_L^0(\alpha_0^L | \mathcal{I}_L) \quad P\text{- and } P_L^0\text{- a.s.} \quad (2.6)$$

Here $g : M_L^\infty \rightarrow \mathbf{R}$ is P -integrable. It holds as well with \mathcal{I}_L replaced by $\mathcal{I}_{L,L'}$. According to the above reference Relation (2.6) can be considered as a conditional version of the inversion formula (1.5).

3 Approximation of P

In this section convergence of the Cesaro mean $t^{-1} \int_0^t P_L^0[T_x \varphi \in B] dx$ will be considered. The limit $Q_L(B)$ is related to the limit $Q_L^0(B)$ of (1.1).

By a generalization of Theorem 3.1 in Nieuwenhuis (1991) (see Section 5 of this reference) it is obvious that

$$\frac{1}{t} \int_0^t 1_B \circ T_x dx \longrightarrow E(1_B | \mathcal{I}_L) \quad P\text{- and } P_L^0\text{-a.s.} \quad (3.1)$$

for all $B \in \mathcal{M}_L^\infty$. The limit equals $P(B)$ if Φ is ergodic. By dominated convergence we have

$$\frac{1}{t} \int_0^t P_L^0[T_x \varphi \in B] dx \longrightarrow E_L^0[E(1_B | \mathcal{I}_L)] =: Q_L(B) \quad (3.2)$$

for all $B \in \mathcal{M}_L^\infty$. Q_L is a probability measure on $(M_L^\infty, \mathcal{M}_L^\infty)$. By Theorem 1.1(ii) and conditioning on \mathcal{I}_L we obtain

$$Q_L(B) = \frac{1}{\lambda(L)} E \left[\frac{1}{\alpha_0^L} E(1_B | \mathcal{I}_L) \right] = \frac{1}{\lambda(L)} E \left[1_B E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \right].$$

Since $E(1/\alpha_0^L | \mathcal{I}_L) > 0$ P -a.s. (see Lemma 2.3(a)),

$$Q_L \sim P \quad \text{and} \quad \frac{dQ_L}{dP} = \frac{1}{\lambda(L)} E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \quad P\text{-a.s.} \quad (3.3)$$

This observation will be used in the proof of the following theorem.

Theorem 3.1 *The convergence in (3.2) holds uniformly for $B \in \mathcal{M}_L^\infty$. If Φ is pseudo- L -ergodic, then $Q_L = P$.*

Proof. Since

$$\frac{1}{t} \int_0^t P_L^0[T_x \varphi \in B] dx = \frac{1}{\lambda(L)t} \int_0^t E \left(\frac{1}{\alpha_0^L} 1_B \circ T_x \circ \vartheta_{0,L} \right) dx,$$

it is sufficient to prove that (3.4) and (3.5) below are satisfied:

$$\sup_{B \in \mathcal{M}_L^\infty} \frac{1}{\lambda(L)t} \left| \int_0^t E \left(\frac{1}{\alpha_0^L} 1_B \circ T_x \circ \vartheta_{0,L} \right) dx - \int_0^t E \left(\frac{1}{\alpha_0^L} 1_B \circ T_x \right) dx \right| \rightarrow 0, \quad (3.4)$$

$$\sup_{B \in \mathcal{M}_L^\infty} \left| \frac{1}{\lambda(L)t} \int_0^t E \left(\frac{1}{\alpha_0^L} 1_B \circ T_x \right) dx - Q_L(B) \right| \rightarrow 0, \quad (3.5)$$

as $t \rightarrow \infty$. First note that

$$\int_0^t 1_B \circ T_x \circ \vartheta_{0,L} dx = \int_{X_0^L}^{X_0^L + t} 1_B \circ T_x dx.$$

Hence,

$$\begin{aligned}
& \frac{1}{\lambda(L)t\alpha_0^L} \left| \int_0^t 1_B \circ T_x \circ \vartheta_{0,L} dx - \int_0^t 1_B \circ T_x dx \right| \leq \\
& \leq \frac{2}{\lambda(L)\alpha_0^L} 1_{\{|X_0^L| > t\}} + \frac{2|X_0^L|}{\lambda(L)t\alpha_0^L} 1_{\{|X_0^L| \leq t\}} \\
& \leq \frac{2}{\lambda(L)\alpha_0^L} 1_{\{|X_0^L| > t\}} + \frac{2}{\lambda(L)t}
\end{aligned}$$

for all $B \in \mathcal{M}_L^\infty$. As a consequence the left-hand side of (3.4) is bounded from above by

$$\frac{2}{\lambda(L)} E \left(\frac{1}{\alpha_0^L} 1_{\{\alpha_0^L > t\}} \right) + \frac{2}{\lambda(L)t} = 2P_L^0[\alpha_0^L > t] + \frac{2}{\lambda(L)t},$$

which tends to 0 as $t \rightarrow \infty$.

Next (3.5). Set

$$\hat{Q}_{t,L}(B) := \frac{1}{\lambda(L)t} \int_0^t E \left(\frac{1}{\alpha_0^L} 1_B \circ T_x \right) dx.$$

Note that $\hat{Q}_{t,L}$ is a probability measure on $(M_L^\infty, \mathcal{M}_L^\infty)$. By (3.1) and dominated convergence we obtain

$$\hat{Q}_{t,L}(B) \longrightarrow E \left[\frac{1}{\lambda(L)\alpha_0^L} E(1_B | \mathcal{I}_L) \right] \text{ as } t \rightarrow \infty. \quad (3.6)$$

Since $E(1_B | \mathcal{I}_L) = E(1_B | \mathcal{I}_L) \circ \vartheta_{0,L}$, the limit in (3.6) equals $Q_L(B)$. By stationarity and Fubini's theorem we have

$$\hat{Q}_{t,L}(B) = \frac{1}{\lambda(L)t} \int_0^t E \left(\frac{1}{\alpha_0^L \circ T_{-x}} 1_B \right) dx = E \left(1_B \frac{1}{\lambda(L)t} \int_0^t \frac{1}{\alpha_0^L \circ T_{-x}} dx \right).$$

So, $\hat{Q}_{t,L}$ is dominated by P and

$$\frac{d\hat{Q}_{t,L}}{dP} = \frac{1}{\lambda(L)t} \int_0^t \frac{1}{\alpha_0^L \circ T_{-x}} dx \quad P\text{-a.s.} \quad (3.7)$$

By an ergodic results similar to (3.1) we obtain

$$\frac{1}{t} \int_0^t \frac{1}{\alpha_0^L \circ T_{-x}} dx \longrightarrow E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \quad P\text{-a.s.}$$

Observe that

$$E \left(\frac{1}{n} \int_0^n \frac{1}{\alpha_0^L \circ T_{-x}} dx \right) = \frac{1}{n} \int_0^n E \left(\frac{1}{\alpha_0^L} \right) dx = \lambda(L) = E \left(E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \right)$$

for all $n \in \mathbf{N}$. Lemma A2.1 in Brandt, Franken and Lisek (1990) ensures that $\left(n^{-1} \int_0^n (\alpha_0^L \circ T_{-x})^{-1} dx \right)_{n \in \mathbf{N}}$ is uniformly P -integrable. By (1.11), (3.7) and (3.3) it is obvious that

$$d(\dot{Q}_{t,L}, Q_L) = E \left| \frac{1}{\lambda(L)t} \int_0^t \frac{1}{\alpha_0^L \circ T_{-x}} dx - \frac{1}{\lambda(L)} E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \right| \rightarrow 0$$

as $\mathbf{N} \ni t \rightarrow \infty$, which proves (3.5) for discrete time-parameter. The transition to continuous time-parameter follows immediately.

The second part of the theorem is a consequence of (3.3), the definition of pseudo- L -ergodicity in (1.9), and Lemma 2.3(a). \square

Note that by stationarity of P and the right-hand part of (2.3),

$$\begin{aligned} Q_L[T_a \varphi \in B] &= \frac{1}{\lambda(L)} E \left[E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) 1_B \circ T_a \right] \\ &= \frac{1}{\lambda(L)} E \left[E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) 1_B \right] = Q_L(B) \end{aligned}$$

for all $B \in \mathcal{M}_L^\infty$. Hence, Q_L is stationary. Since $Q_L = P$ and $Q_L^0 = P_L^0$ (see (1.10)) provided that Φ is pseudo- L -ergodic, one might wonder if Q_L^0 is the Palm distribution with respect to L associated with Q_L .

To prove that this is usually not the case, let \tilde{Q}_L^0 be this Palm distribution associated with Q_L and let $\tilde{\lambda}(L)$ be the intensity of Q_L . Recall the definition of N_1^L in Section 2. By (3.3), conditioning on \mathcal{I}_L , Theorem 1.1, and Lemma 2.3 we have

$$\tilde{\lambda}(L) = E_{Q_L} N_1^L = \frac{1}{\lambda(L)} E \left(\frac{1}{\alpha_0^L} E(N_1^L | \mathcal{I}_L) \right) = E_L^0(E(N_1^L | \mathcal{I}_L)) = E_L^0(1/E_L^0(\alpha_0^L | \mathcal{I}_L)).$$

Since (cf. (1.3) and (3.3))

$$\begin{aligned}
\tilde{Q}_L^0(B) &= \frac{1}{\bar{\lambda}(L)} E_{Q_L} \sum_{i=1}^{N_L^L} 1_B \circ \vartheta_{i,L} \\
&= \frac{1}{\bar{\lambda}(L)\lambda(L)} E \left[\sum_{i=1}^{N_L^L} 1_B \circ \vartheta_{i,L} \cdot E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \right] \\
&= \frac{1}{\bar{\lambda}(L)\lambda(L)} E \left[\sum_{i=1}^{N_L^L} \left(1_B E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \right) \circ \vartheta_{i,L} \right] \\
&= \frac{1}{\bar{\lambda}(L)} E_L^0 \left(1_B E \left(\frac{1}{\alpha_0^L} | \mathcal{I}_L \right) \right)
\end{aligned}$$

for all $B \in \mathcal{M}_L^\infty$, we obtain

$$\tilde{Q}_L^0(B) = \frac{E_L^0(1_B/E_L^0(\alpha_0^L|\mathcal{I}_L))}{E_L^0(1/E_L^0(\alpha_0^L|\mathcal{I}_L))}, \quad B \in \mathcal{M}_L^\infty. \quad (3.8)$$

Consequently,

$$\tilde{Q}_L^0 \sim P_L^0 \quad \text{and} \quad \frac{d\tilde{Q}_L^0}{dP_L^0} = \frac{1/E_L^0(\alpha_0^L|\mathcal{I}_L)}{E_L^0(1/E_L^0(\alpha_0^L|\mathcal{I}_L))}. \quad (3.9)$$

Hence (cf. Theorem 1.2),

$$\frac{d\tilde{Q}_L^0}{dQ_L^0} = \frac{d\tilde{Q}_L^0}{dP_L^0} \frac{dP_L^0}{dQ_L^0} = \frac{1/(E_L^0(\alpha_0^L|\mathcal{I}_L))^2}{\lambda(L)E_L^0(1/E_L^0(\alpha_0^L|\mathcal{I}_L))}$$

and

$$\tilde{Q}_L^0 = Q_L^0 \quad \text{iff} \quad \Phi \text{ is pseudo-}L\text{-ergodic.}$$

This last result ensures that Q_L^0 is the Palm distribution with respect to L associated with Q_L iff Φ is pseudo- L -ergodic.

4 Approximation of P_L^0 starting from $P_{L'}^0$

Let $L, L' \in \text{Bor } K$ be such that $L \cap L' = \emptyset$ and $P(M_{L,L'}^\infty) = 1$. In this section our emphasis is on the Palm distributions P_L^0 and $P_{L'}^0$, now considered as probability measures on $(M_{L,L'}^\infty, \mathcal{M}_{L,L'}^\infty)$. Relation (1.10) expresses uniform approximation of P_L^0 , starting from P . It holds as well with $P_{L'}^0$, instead of P .

Since the restriction of $E_L^0(1_B | \mathcal{I}_L)$ to $M_{L,L'}^\infty$ equals $E_L^0(1_B | \mathcal{I}_{L,L'})$ P_L^0 - and P -a.s., the following cross ergodic theorem holds:

$$\frac{1}{n} \sum_{i=1}^n 1_B \circ \vartheta_{i,L} \longrightarrow E_L^0(1_B | \mathcal{I}_{L,L'}) \quad P\text{-a.s.}, \quad (4.1)$$

$B \in \mathcal{M}_{L,L'}^\infty$. Our starting point is now $P_{L'}^0$. So, we are interested in the probability measure $Q_{L,L'}$ defined by

$$Q_{L,L'}(B) := E_{L'}^0[E_L^0(1_B | \mathcal{I}_{L,L'})], \quad B \in \mathcal{M}_{L,L'}^\infty. \quad (4.2)$$

Since $\mathcal{I}_{L,L'} = \mathcal{I}_{L',L}$ (cf. Lemma 2.2), it follows that

$$P_{0,L'}(A) = E(1_A \circ \vartheta_{0,L'}) = E(1_A) = E(1_A \circ \vartheta_{0,L}) = P_{0,L}(A)$$

for all $A \in \mathcal{I}_{L,L'}$. Set $M^0 := M_L^0 \cap M_{L'}^\infty$ and $A_0 := [E_L^0(1_{M^0} | \mathcal{I}_{L,L'}) = 1]$. Since $P_L^0(A_0) = 1$, $A_0 \in \mathcal{I}_{L,L'}$, and $P_L^0 \sim P_{0,L}$, we obtain: $P_{L'}^0(A_0) = 1$. Consequently, $Q_{L,L'}(M^0) = 1$.

Using the Radon-Nikodym derivatives $dP_{L'}^0/dP_{0,L'}$ and $dP_{0,L}/dP_L^0$, cf. Theorem 1.1, we obtain

$$\begin{aligned} Q_{L,L'}(B) &= \frac{1}{\lambda(L')} E \left[\frac{1}{\alpha_0^{L'}} E_L^0(1_B | \mathcal{I}_{L,L'}) \right] \\ &= \frac{1}{\lambda(L')} E \left[E \left(\frac{1}{\alpha_0^{L'}} | \mathcal{I}_{L,L'} \right) E_L^0(1_B | \mathcal{I}_{L,L'}) \right] \\ &= \frac{\lambda(L)}{\lambda(L')} E_L^0 \left[\alpha_0^L E \left(\frac{1}{\alpha_0^{L'}} | \mathcal{I}_{L,L'} \right) E_L^0(1_B | \mathcal{I}_{L,L'}) \right] \\ &= \frac{\lambda(L)}{\lambda(L')} E_L^0 \left[E_L^0(1_B E_L^0(\alpha_0^L | \mathcal{I}_{L,L'}) E \left(\frac{1}{\alpha_0^{L'}} | \mathcal{I}_{L,L'} \right) | \mathcal{I}_{L,L'} \right] \end{aligned}$$

$$= \frac{\lambda(L)}{\lambda(L')} E_L^0 \left[1_B \frac{E_L^0(\alpha_0^L | \mathcal{I}_{L,L'})}{E_{L'}^0(\alpha_0^{L'} | \mathcal{I}_{L,L'})} \right],$$

$B \in \mathcal{M}_{L,L'}^\infty$. In the second and fourth equality we conditioned on $\mathcal{I}_{L,L'}$; the last equality is a consequence of Lemma 2.3(b). Consequently, on $(M_{L,L'}^\infty, \mathcal{M}_{L,L'}^\infty)$ we have:

$$Q_{L,L'} \sim P_L^0 \quad \text{and} \quad \frac{dQ_{L,L'}}{dP_L^0} = \frac{\lambda(L) E_L^0(\alpha_0^L | \mathcal{I}_{L,L'})}{\lambda(L') E_{L'}^0(\alpha_0^{L'} | \mathcal{I}_{L,L'})}. \quad (4.3)$$

This observation is essential for the following theorem.

Theorem 4.1

- (a) Relation (4.1) holds as well with $P_{L'}^0$ instead of P .
 (b) The convergence

$$\frac{1}{n} \sum_{i=1}^n P_{L'}^0[\vartheta_{i,L} \varphi \in B] \longrightarrow Q_{L,L'}(B) \quad (4.4)$$

is uniform for $B \in \mathcal{M}_{L,L'}^\infty$.

- (c) If Φ is pseudo- L -ergodic and pseudo- L' -ergodic, then $Q_{L,L'}$ equals $P_{L'}^0$.

Proof. Let $\nu : M_{L,L'}^\infty \rightarrow \mathbf{N}_0$ be defined by $\nu(\varphi) := \varphi_L(X_0^{L'}(\varphi), 0]$, $\varphi \in M_{L,L'}^\infty$. Since $\vartheta_{i,L} \circ \vartheta_{0,L'} \varphi = \varphi(X_{i-\nu(\varphi)}^L(\varphi) + \cdot)$, we obtain

$$\begin{aligned} & P_{0,L'} \left[\frac{1}{n} \sum_{i=1}^n 1_B \circ \vartheta_{i,L} \rightarrow E_L^0(1_B | \mathcal{I}_{L,L'}) \right] \\ &= P \left[\frac{1}{n} \sum_{i=1}^n 1_B \circ \vartheta_{i-\nu,L} \rightarrow E_L^0(1_B | \mathcal{I}_{L,L'}) \right] \\ &= P \left[\frac{1}{n} \sum_{j=1-\nu}^{n-\nu} 1_B \circ \vartheta_{j,L} \rightarrow E_L^0(1_B | \mathcal{I}_{L,L'}) \right], \end{aligned}$$

$B \in \mathcal{M}_{L,L'}^\infty$. The last probability equals 1 since (4.1) remains valid if in the summation n is replaced by $n - \nu$. Since $P_{L'}^0 \sim P_{0,L'}$, part (a) follows immediately.

By dominated convergence it is obvious that (4.4) holds for all $B \in \mathcal{M}_{L,L'}^\infty$; only the uniformity of the convergence remains to prove. Define

$$R_n(B, \varphi) := \frac{1}{\lambda(L')\alpha_0^{L'}(\varphi)} \frac{1}{n} \sum_{j=1-\nu(\varphi)}^{n-\nu(\varphi)} 1_B \circ \vartheta_{j,L}(\varphi),$$

$$S_n(B, \varphi) := \frac{1}{\lambda(L')\alpha_0^{L'}(\varphi)} \frac{1}{n} \sum_{j=1}^n 1_B \circ \vartheta_{j,L}(\varphi),$$

for $\varphi \in M_{L,L'}^\infty$ and $B \in \mathcal{M}_{L,L'}^\infty$. Note that the left-hand side in (4.4) equals $ER_n(B)$. Since

$$|R_n(B, \varphi) - S_n(B, \varphi)| \leq \frac{2}{\lambda(L')\alpha_0^{L'}(\varphi)} 1_{[\nu \geq n]}(\varphi) + \frac{2\nu(\varphi)}{n\lambda(L')\alpha_0^{L'}(\varphi)} 1_{[\nu < n]}(\varphi)$$

for all $\varphi \in M_{L,L'}^\infty$, we obtain:

$$E|R_n(B) - S_n(B)| \leq \frac{2}{\lambda(L')} E \left[\frac{1}{\alpha_0^{L'}} 1_{[\nu \geq n]} \right] + \frac{2}{n\lambda(L')} E \left[\frac{\nu}{\alpha_0^{L'}} \right],$$

uniformly for $B \in \mathcal{M}_{L,L'}^\infty$. This upper bound tends to 0 as $n \rightarrow \infty$. Hence, it is sufficient to prove that

$$\sup_{B \in \mathcal{M}_{L,L'}^\infty} |ES_n(B) - Q_{L,L'}(B)| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (4.5)$$

For this we define probability measures $\mu_{i,L}$ as follows

$$\mu_{i,L}(B) := E \left[\frac{1}{\lambda(L')\alpha_0^{L'}} 1_B \circ \vartheta_{i,L} \right], \quad B \in \mathcal{M}_{L,L'}^\infty.$$

By (1.5), Fubini's theorem, and (1.4) we obtain:

$$\begin{aligned} \mu_{i,L}(B) &= \frac{\lambda(L)}{\lambda(L')} \int_0^\infty E_L^0 \left[\frac{1}{\alpha_0^{L'} \circ T_u} (1_B \circ \vartheta_{i,L} \circ T_u) 1_{[\alpha_0^L > u]} \right] du \\ &= \frac{\lambda(L)}{\lambda(L')} E_L^0 \left[\int_0^{\alpha_0^L} \frac{1}{\alpha_0^{L'} \circ T_u} 1_B \circ \vartheta_{i,L} du \right] \\ &= \frac{\lambda(L)}{\lambda(L')} E_L^0(1_B \eta_i), \end{aligned}$$

where

$$\eta_i = \int_0^{\alpha_L^L} \frac{1}{\alpha_0^{L'} \circ T_u \circ \vartheta_{-i,L}} du.$$

Hence, for all i and $n \in \mathbf{N}$ we have

$$\mu_{i,L} \ll P_L^0 \quad \text{and} \quad \frac{d\mu_{i,L}}{dP_L^0} = \frac{\lambda(L)}{\lambda(L')} \eta_i, \quad (4.6)$$

$$ES_n \ll P_L^0 \quad \text{and} \quad \frac{dES_n}{dP_L^0} = \frac{\lambda(L)}{\lambda(L')} \frac{1}{n} \sum_{i=1}^n \eta_i. \quad (4.7)$$

By (4.3), (4.7), and the definition of d just above (1.11) we obtain

$$d(ES_n, Q_{L,L'}) = \frac{\lambda(L)}{\lambda(L')} E_L^0 \left| \frac{1}{n} \sum_{i=1}^n \eta_i - \frac{E_L^0(\alpha_0^L | \mathcal{I}_{L,L'})}{E_{L'}^0(\alpha_0^{L'} | \mathcal{I}_{L,L'})} \right|. \quad (4.8)$$

Note that (η_i) is P_L^0 -stationary. So,

$$\frac{1}{n} \sum_{i=1}^n \eta_i \longrightarrow E_L^0 \left(\int_0^{\alpha_0^L} \frac{1}{\alpha_0^{L'} \circ T_u} du | \mathcal{I}_{L,L'} \right) \quad P_L^0\text{-a.s.} \quad (4.9)$$

By (2.6) and Lemma 2.3(b) the limit in (4.9) is equal to

$$E \left(\frac{1}{\alpha_0^{L'}} | \mathcal{I}_{L,L'} \right) E_L^0(\alpha_0^L | \mathcal{I}_{L,L'}) = \frac{E_L^0(\alpha_0^L | \mathcal{I}_{L,L'})}{E_{L'}^0(\alpha_0^{L'} | \mathcal{I}_{L,L'})} \quad P_L^0\text{-a.s.} \quad (4.10)$$

Since the sequence $(\sum_{i=1}^n \eta_i/n)$ is uniformly P_L^0 -integrable (cf., e.g., Brandt, Franken and Lisek (1990; L.A2.1)), we obtain by (4.8)-(4.10) that

$$d(ES_n, Q_{L,L'}) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

see also Brémaud (1981; Th.T26). This proves (4.5) and hence part (b).

Part (c) is an immediate consequence of (4.3) and Lemma 2.3. \square

References

- Brandt, A., P. Franken and B. Lisek (1990). *Stationary Stochastic Models*, Wiley, New York.
- Brémaud, P. (1981). *Point Processes and Queues*, Springer, New York.
- Franken, P., D. König, U. Arndt and V. Schmidt (1982). *Queues and Point Processes*, Wiley, New York.
- Kallenberg, O. (1983/86). *Random Measures*, 3rd and 4th editions, Akademie-Verlag and Academic Press, Berlin and London.
- König, D. and V. Schmidt (1986). Limit theorems for single-server feedback queues controlled by a general class of marked point processes, *Theory Prob. Appl.* **30**, 712-719.
- Konstantopoulos, P. and J. Walrand (1988). On the weak convergence of stochastic processes with embedded point processes, *Adv. Appl. Prob.* **20**, 473-475.
- Matthes, K., J. Kerstan and J. Mecke (1978). *Infinitely Divisible Point Processes*, Wiley, New York.
- Nieuwenhuis, G. (1989). Equivalence of functional limit theorems for stationary point processes and their Palm distributions, *Probability Theory and Related Fields* **81**, 593-608.
- Nieuwenhuis, G. (1991). Bridging the gap between a stationary point process and its Palm distribution, Research Memorandum FEW 502, Department of Economics, University of Tilburg.

IN 1991 REEDS VERSCHENEN

- 466 Prof.Dr. Th.C.M.J. van de Klundert - Prof.Dr. A.B.T.M. van Schaik
Economische groei in Nederland in een internationaal perspectief
- 467 Dr. Sylvester C.W. Eijffinger
The convergence of monetary policy - Germany and France as an example
- 468 E. Nijssen
Strategisch gedrag, planning en prestatie. Een inductieve studie binnen de computerbranche
- 469 Anne van den Nouweland, Peter Borm, Guillermo Owen and Stef Tijs
Cost allocation and communication
- 470 Drs. J. Grazell en Drs. C.H. Veld
Motieven voor de uitgifte van converteerbare obligatieleningen en warrant-obligatieleningen: een agency-theoretische benadering
- 471 P.C. van Batenburg, J. Kriens, W.M. Lammerts van Bueren and R.H. Veenstra
Audit Assurance Model and Bayesian Discovery Sampling
- 472 Marcel Kerkhofs
Identification and Estimation of Household Production Models
- 473 Robert P. Gilles, Guillermo Owen, René van den Brink
Games with Permission Structures: The Conjunctive Approach
- 474 Jack P.C. Kleijnen
Sensitivity Analysis of Simulation Experiments: Tutorial on Regression Analysis and Statistical Design
- 475 C.P.M. van Hoesel
An $O(n \log n)$ algorithm for the two-machine flow shop problem with controllable machine speeds
- 476 Stephan G. Vanneste
A Markov Model for Opportunity Maintenance
- 477 F.A. van der Duyn Schouten, M.J.G. van Eijs, R.M.J. Heuts
Coordinated replenishment systems with discount opportunities
- 478 A. van den Nouweland, J. Potters, S. Tijs and J. Zarzuelo
Cores and related solution concepts for multi-choice games
- 479 Drs. C.H. Veld
Warrant pricing: a review of theoretical and empirical research
- 480 E. Nijssen
De Miles and Snow-typologie: Een exploratieve studie in de meubelbranche
- 481 Harry G. Barkema
Are managers indeed motivated by their bonuses?

- 482 Jacob C. Engwerda, André C.M. Ran, Arie L. Rijkeboer
Necessary and sufficient conditions for the existence of a positive definite solution of the matrix equation $X + A^T X^{-1} A = I$
- 483 Peter M. Kort
A dynamic model of the firm with uncertain earnings and adjustment costs
- 484 Raymond H.J.M. Gradus, Peter M. Kort
Optimal taxation on profit and pollution within a macroeconomic framework
- 485 René van den Brink, Robert P. Gilles
Axiomatizations of the Conjunctive Permission Value for Games with Permission Structures
- 486 A.E. Brouwer & W.H. Haemers
The Gewirtz graph - an exercise in the theory of graph spectra
- 487 Pim Adang, Bertrand Melenberg
Intratemporal uncertainty in the multi-good life cycle consumption model: motivation and application
- 488 J.H.J. Roemen
The long term elasticity of the milk supply with respect to the milk price in the Netherlands in the period 1969-1984
- 489 Herbert Hamers
The Shapley-Entrance Game
- 490 Rezaul Kabir and Theo Vermaelen
Insider trading restrictions and the stock market
- 491 Piet A. Verheyen
The economic explanation of the jump of the co-state variable
- 492 Drs. F.L.J.W. Manders en Dr. J.A.C. de Haan
De organisatorische aspecten bij systeemontwikkeling een beschouwing op besturing en verandering
- 493 Paul C. van Batenburg and J. Kriens
Applications of statistical methods and techniques to auditing and accounting
- 494 Ruud T. Frambach
The diffusion of innovations: the influence of supply-side factors
- 495 J.H.J. Roemen
A decision rule for the (des)investments in the dairy cow stock
- 496 Hans Kremers and Dolf Talman
An SLSPP-algorithm to compute an equilibrium in an economy with linear production technologies

- 497 L.W.G. Strijbosch and R.M.J. Heuts
Investigating several alternatives for estimating the compound lead
time demand in an (s,Q) inventory model
- 498 Bert Bettonvil and Jack P.C. Kleijnen
Identifying the important factors in simulation models with many
factors
- 499 Drs. H.C.A. Roest, Drs. F.L. Tijssen
Beheersing van het kwaliteitsperceptieproces bij diensten door middel
van keurmerken
- 500 B.B. van der Genugten
Density of the F-statistic in the linear model with arbitrarily
normal distributed errors
- 501 Harry Barkema and Sytse Douma
The direction, mode and location of corporate expansions
- 502 Gert Nieuwenhuis
Bridging the gap between a stationary point process and its Palm
distribution
- 503 Chris Veld
Motives for the use of equity-warrants by Dutch companies
- 504 Pieter K. Jagersma
Een etiologie van horizontale internationale ondernemingsexpansie
- 505 B. Kaper
On M-functions and their application to input-output models
- 506 A.B.T.M. van Schaik
Productiviteit en Arbeidsparticipatie
- 507 Peter Borm, Anne van den Nouweland and Stef Tijs
Cooperation and communication restrictions: a survey
- 508 Willy Spanjers, Robert P. Gilles, Pieter H.M. Ruys
Hierarchical trade and downstream information
- 509 Martijn P. Tummers
The Effect of Systematic Misperception of Income on the Subjective
Poverty Line
- 510 A.G. de Kok
Basics of Inventory Management: Part 1
Renewal theoretic background
- 511 J.P.C. Blanc, F.A. van der Duyn Schouten, B. Pourbabai
Optimizing flow rates in a queueing network with side constraints
- 512 R. Peeters
On Coloring j-Unit Sphere Graphs

- 513 Drs. J. Dagevos, Drs. L. Oerlemans, Dr. F. Boekema
Regional economic policy, economic technological innovation and networks
- 514 Erwin van der Krabben
Het functioneren van stedelijke onroerend-goed-markten in Nederland - een theoretisch kader
- 515 Drs. E. Schaling
European central bank independence and inflation persistence
- 516 Peter M. Kort
Optimal abatement policies within a stochastic dynamic model of the firm
- 517 Pim Adang
Expenditure versus consumption in the multi-good life cycle consumption model
- 518 Pim Adang
Large, infrequent consumption in the multi-good life cycle consumption model
- 519 Raymond Gradus, Sjak Smulders
Pollution and Endogenous Growth
- 520 Raymond Gradus en Hugo Keuzenkamp
Arbeidsongeschiktheid, subjectief ziektegevoel en collectief belang
- 521 A.G. de Kok
Basics of inventory management: Part 2
The (R,S)-model
- 522 A.G. de Kok
Basics of inventory management: Part 3
The (b,Q)-model
- 523 A.G. de Kok
Basics of inventory management: Part 4
The (s,S)-model
- 524 A.G. de Kok
Basics of inventory management: Part 5
The (R,b,Q)-model
- 525 A.G. de Kok
Basics of inventory management: Part 6
The (R,s,S)-model
- 526 Rob de Groof and Martin van Tuijl
Financial integration and fiscal policy in interdependent two-sector economies with real and nominal wage rigidity

- 527 A.G.M. van Eijs, M.J.G. van Eijs, R.M.J. Heuts
Gecoördineerde bestelsystemen
een management-georiënteerde benadering
- 528 M.J.G. van Eijs
Multi-item inventory systems with joint ordering and transportation
decisions
- 529 Stephan G. Vanneste
Maintenance optimization of a production system with buffercapacity
- 530 Michel R.R. van Bremen, Jeroen C.G. Zijlstra
Het stochastische variantie optiewaarderingsmodel
- 531 Willy Spanjers
Arbitrage and Walrasian Equilibrium in Economies with Limited Infor-
mation

IN 1992 REEDS VERSCHENEN

- 532 F.G. van den Heuvel en M.R.M. Turlings
Privatisering van arbeidsongeschiktheidsregelingen
Refereed by Prof.Dr. H. Verbon
- 533 J.C. Engwerda, L.G. van Willigenburg
LQ-control of sampled continuous-time systems
Refereed by Prof.dr. J.M. Schumacher
- 534 J.C. Engwerda, A.C.M. Ran & A.L. Rijkeboer
Necessary and sufficient conditions for the existence of a positive definite solution of the matrix equation $X + A^*X^{-1}A = Q$.
Refereed by Prof.dr. J.M. Schumacher
- 535 Jacob C. Engwerda
The indefinite LQ-problem: the finite planning horizon case
Refereed by Prof.dr. J.M. Schumacher
- 536 Gert-Jan Otten, Peter Borm, Ton Storcken, Stef Tijs
Effectivity functions and associated claim game correspondences
Refereed by Prof.dr. P.H.M. Ruys
- 537 Jack P.C. Kleijnen, Gustav A. Alink
Validation of simulation models: mine-hunting case-study
Refereed by Prof.dr.ir. C.A.T. Takkenberg
- 538 V. Feltkamp and A. van den Nouweland
Controlled Communication Networks
Refereed by Prof.dr. S.H. Tijs
- 539 A. van Schaik
Productivity, Labour Force Participation and the Solow Growth Model
Refereed by Prof.dr. Th.C.M.J. van de Klundert
- 540 J.J.G. Lemmen and S.C.W. Eijffinger
The Degree of Financial Integration in the European Community
Refereed by Prof.dr. A.B.T.M. van Schaik
- 541 J. Bell, P.K. Jagersma
Internationale Joint Ventures
Refereed by Prof.dr. H.G. Barkema
- 542 Jack P.C. Kleijnen
Verification and validation of simulation models
Refereed by Prof.dr.ir. C.A.T. Takkenberg

Bibliotheek K. U. Brabant



17 000 01087535 0